

# Australia & New Zealand Pancreas Transplant Registry Report 1984-2007

---



This report is a compilation of data provided by the five current Pancreas transplant units in Australia and New Zealand: Auckland Renal Transplant Groups, New Zealand; Monash Medical Centre, Clayton, Victoria; Royal Prince Alfred hospital Camperdown, Diabetes Transplant unit, Randwick, and National Pancreas Transplant Unit, Westmead Hospital NSW Australia. The registry is funded in part by a grant from the Commonwealth Department of Health and Ageing.

*Southern Health*

Inquiries or comments should be directed  
to the secretariat at:  
National Pancreas Transplant Unit  
Westmead NSW 2145  
Ph: 61 2 9845 6962 > Fax: 61 2 9633 9351  
Email: paul\_robertson@wsahs.nsw.gov.au



**Edited by**

Tracy Brislane, Jeremy Chapman & Paul Robertson

**Participating centers**

|                   |   |
|-------------------|---|
| Dr Helen Pilmore  | Auckland Renal Transplant Group; Auckland City Hospital |
| Ms Elaine Kennedy | Monash Medical Centre                                   |
| Mr Paul Robertson | Renal and Urology, Westmead Hospital                    |
| Ms Jane Mawson    | Royal Prince Alfred Hospital, Camperdown                |
|                   | Prince of Wales Hospital, Randwick                      |

**National Pancreas Transplant Registry**

|                         |  |
|-------------------------|--|
| Dr Henry Pleass         | Director, National Pancreas Transplant Unit                  |
| Prof Jeremy Chapman     | Director of Acute Interventional Medicine, Westmead Hospital |
| A/Prof Philip O'Connell | Physician  |
| Dr Brian Nankivell      | Physician  |
| Prof Richard Allen      | Surgeon  |
| Dr Howard Lau           | Surgeon  |
| Mr Paul Robertson       | Transplant Co-ordinator                                      |
| Mr Lincoln Dealtry      | Living related transplant Co-ordinator                       |
| Ms Kathy Kable          | Clinical Nurse Consultant                                    |
| Dr Wayne Hawthorne      | Senior Research Fellow                                       |
| Ms Tracy Brislane       | Data Manager   |
| Ms Penny Murie          | Clinical Research Co-ordinator                               |
| Ms Julie McKelvey       | Registry clerk   |

March 2008



## Table of Content

*Table of Content*..... 1

*Summary*..... 7

*Introduction*..... 4

*Analysis*..... 4

*Number of transplants* ..... 5

*Pancreas Transplant by Category* ..... 6

*By Pancreas duct management*..... 7

*Preservation time*..... 8

*Demographics*..... 8

*Gender*..... 8

*Age groups*..... 9

*Smoking* ..... 9

*Alcohol status in donor* ..... 10

*Islet transplant*..... 10

*Death*..... 11

*Reasons for graft failure*..... 11

*Re-transplant*..... 12

*Waiting time* ..... 12

*Organ donation and exchange* ..... 13

*Bibliography* ..... 16

## Summary

- ▲ 412 Pancreas transplants have been performed in Australia and New Zealand (ANZ) in 1984-2007 including Islet transplants.
- ▲ In 2007, 31 transplants were performed: Auckland (1), Monash (9), Prince of Wales (0), Westmead (21). Transplant by category: SPK (26), PTA (1), PAK (2) and ITA (2).

### International

- ▲ ANZ constituted of 8% (24 out of 433) to the non-US transplant activity in 2002.
- ▲ One-year SPK Patient survival was slightly higher in ANZ (97%) than in US (95%) and non-US (96%) in 1996-2002.
- ▲ ANZ had similar proportion of SPK to non-US (90%) but less PAK and PTA.
- ▲ ANZ compared favourably with international data on the 1-year Pancreas survival, Patient survival, and Kidney survival.

### SPK Patient survival (Figure i)

- ▲ The 1-year, 3-year and 5-year SPK Patient survival was 96%, 95% and 94% in 1984-2007.
- ▲ Recipients had significantly 4 fold lower in Instantaneous relative risk (hazard rate, HR) of death if the operation was performed after 1994.

### SPK Kidney graft survival (Figure i)

- ▲ 1-year, 3- year and 5-year was 97%, 95% and 94%.

#### 1-year

- ▲ 96% in 1984-1993 and 97% 1994-2004.
- ▲ 98% for BD and 97% for ED.
- ▲ 97% for the recipients aged <45 and 94% ≥45.

### SPK Pancreas graft survival (Figure i)

- ▲ 1-year, 3-year and 5-year Pancreas survival was 86%, 84% and 80% respectively in 1984-2007.

#### 1- year

- ▲ 78% in 1984-1993 and 87% in 1994-2004.

- ▲ 86% for BD and 90% for ED.

- ▲ 86% for the recipients aged <45 and 85% for those ≥45.

### Islet transplant

Twenty-one Islet transplant procedures have been performed in nine patients since 2002. 2 were performed in 2007 in Westmead.

### Others

- ▲ The most common known causes of death was Cardio/Cerebrovascular event (39%)
- ▲ 39% of kidney and pancreas graft failure was due to thrombosis.
- ▲ Eight recipients had a second transplant: SPK (4), SPK followed by PTA (1), PAK (3).

### Glossary

- ▲ SPK - Simultaneous Kidney Pancreas Transplant
- ▲ PTA - Pancreas Transplant Alone
- ▲ PAK - Pancreas after Kidney Transplant
- ▲ ITA - Islet Transplant Alone
- ▲ BD - Bladder Drained Pancreas
- ▲ ED – Enteric Drained Pancreas

Figure i. Patient and Pancreas graft survival in Australia and New Zealand, 1984-2007





Figure ii Patient and Graft Survival from ANZDATA between 1997 - 2006



Source: ANZDATA Registry Report 2007

## Introduction


The Australia and New Zealand (ANZ) Pancreas transplant data were obtained from the transplant centers in Auckland Renal Transplant Group (Auckland), Monash Medical Center (Monash), Royal Prince Alfred Hospital , (Camperdown) and National Pancreas Transplant Unit at Westmead Hospital (Westmead). Auckland commenced in 1998. Islets were first performed in Westmead in 2002, and subsequently performed also at Prince of Wales hospital.

There have been 412 pancreas transplants in Australia and New Zealand (ANZ) between the dates of November 1984 to December 2007 including Islet transplants. No activity was recorded in 1985 and 1986. Of these, nine patients have had a second transplant.

## Analysis

The functioning pancreas grafts are defined as insulin independent recipients. Kidney grafts are defined as functioning if recipients are dialysis free. All causes of death are included in the survival analysis. Patients receiving a second transplant after failure of the first graft are censored for Patient survival for the first graft at the date of the second graft.

The aims of this report are to:

- outline all the transplant activity in 1984-2007
- compare the ANZ data to US and non US (noted by  in the text )

The distribution of Patient survival, Kidney graft survival and Pancreas graft survival was analysed by two periods (1984-1993 and 1994-2007), pancreas duct management (Bladder drained [BD] and Enteric drained [ED]) and two age groups (below 45 and above 45) in Simultaneous pancreas and kidney (SPK) recipients. A brief discussion was made on duct management, causes of deaths, donor age, causes of graft failure, re-transplant, waiting list and number of organ donation.

Kaplan-Meier survival curves were used to illustrate the survival distributions. Cox regression models were used to estimate the Instantaneous relative risks (hazard ratios, HR) and their 95% confidence intervals. The HR quantifies differences in survival between groups. The statistical software package, SPSS® for Windows Release 13.0 was used for all analyses.

## Number of transplants

Figure 1 illustrates the number of transplants in ANZ between 1984 and 2007. Eighty six percent of the transplants were performed after 1994 (353 out of 412) including Islet transplants. Between 1984-2007, the transplants were performed in Westmead (281 out of 406), Monash (99), Auckland (23), RPA (1), RMH (1), QEH (1) and POW (6). In 2007, 31 transplants were performed: Auckland (1), Monash (9), and Westmead (21).




Figure 2. Number of Pancreas transplants by centre in Australia and New Zealand, 1984-2007

## Pancreas Transplant by Category

Figure 2 shows the pancreas transplants by category: Simultaneous pancreas-kidney transplant (SPK), Pancreas after kidney (PAK), Pancreas transplant alone (PTA) and Islets transplant (ITA). SPK is the major pancreas transplants by category in ANZ at 91% (374 out of 412) in 1984-2007 followed by ITA (21), PAK (9), PTA (7) and one Pancreas/Liver and Kidney which was performed in 2005.

**Pancreas transplant by Category 1984 - 2007**




Figure 2. Pancreas transplant by category in Australia and New Zealand, 1984-2007



The proportion of all transplants that were SPK was over 90% in both ANZ (374 out of 391, excluding twenty one ITA) and non-US (4,336 out of 4,756) but less in US (79%, 11,505 out of 14,605) in 1984-2004 (Figure 3). In 2002, ANZ (4%, 0%) had less PAK and PTA compared to non-US (11%, 6%) and US (26%, 9%).






Figure 3. Pancreas transplant by category (%)

**By Pancreas duct management**

ED was introduced in ANZ during 2001. Figure 5 illustrates the number of transplant by duct management. In 2007, all transplants activities were performed with ED but four.

**Pancreas Duct Management - 1987 - 2007**




Figure 4. Pancreas duct management in ANZ, 1984-2007

## Preservation time

The mean ischaemic time was 10-11 hours (standard deviation: 4 hours) for both kidney and pancreas.




Figure 5. Ischaemic time for Kidney and Pancreas graft, 1984-2007

## Demographics

### Gender

63% of donors were male compared to 50% in the recipients (Figure 6).




Figure 6. Gender in Donor and recipients, 1984-2007

### Age groups

Figure illustrates the distribution of six age groups in both donor and recipients. The age range for the donors was between 4 to 61 years and 20 to 60 in the recipients. The median age for donors was 24 years (standard deviation: 10.2) and 37 years (standard deviation: 7.00) for the recipients. Majority of the donors and recipients were aged less than 40.




Figure 7. Age groups in Donor and recipients, 1984-2007

### Smoking

Majority of the recipients had never smoked (57%) compared to 47% in the recipients. Only 4 percent of the recipients were current smokers.




Figure 8. Smoking status in Donor and recipients, 1984-2007

### Alcohol status in donor

Seventy percent of the donors had never consumed alcohol (> than 40g/day) (Figure9).




Figure 9 Alcohol consumption in Donor: 1984-2007

### Islet transplant

Since 2002, twenty one Islet transplant procedures have been performed in 9 patients with 100% patient survival. Figure 18 shows the Islet recipient by age group. The mean age of Islet recipients at transplant was 43 (standard deviation: 7.0). There were an equal proportion of recipients in both sexes. The average length of Type I Diabetes to transplant was 22 years, (standard deviation: 5.6 years).

**Islet Transplants - byAge %: 2002 - 2007**





Figure 10. Islet recipient by age groups at transplant, 2002-2006

 The number of Islet transplant worldwide in 2000 was 280

## Death

Figure 11 shows the causes of death after transplant for recipients in 1984-2007. The common known causes of death were Cardio/Cerebrovascular event 39% and Infection at 17%.

**Causes of Death 1984 - 2007**




Figure 11. Number of known causes of death after transplant, 1984-2006

## Reasons for graft failure

57 percent of pancreas graft failure were due to thrombosis in 1984-2007 (Figure 12 ). The major cause of kidney graft failure was rejection at 37%.

**Graft Failure 1984 - 2007**





Figure 12. Reasons for graft failure

 Thrombosis was also the most common reason for graft failure in US, 1999-2003 (60%).

## Re-transplant

Nine recipients had a second transplant. Four pancreas recipients had a second SPK and one had a PTA and three had a PAK.

## Waiting time

In December 2006, the number of recipients on the active Kidney Pancreas Transplant waiting list was 37. The number of patients on Islet Transplant Waiting list was 6 (The average waiting time for patients transplanted with cadaver pancreas was 1.9 years (33 patients) and 1.4 years (4 patients) for islets.




Figure 13. Waiting list for Pancreas transplant in Australia

Source: ANZOD registry Report 2007

## Organ donation and exchange

During 1994 to 2007, the average number of donated cadaver pancreases and kidneys was 43 and 362 respectively (Figure 14).




Figure 14. Annual number of donated kidney, pancreas and islets in Australia, 1994-2007

Source: ANZOD Registry Report 2008

Figure 15 shows the number of pancreas transplant retrieved and transplant by Australia States and New Zealand in 1989-2007.

**Pancreas transplant retrievals 1989 - 2007**




Figure 15. Number of pancreas transplanted and retrieved by Australian States and New Zealand, 1989-2006

Source: ANZOD Registry Report 2007



The donation of pancreas from States and New Zealand in 2005-2006 is illustrated in Figure 16.




Figure 16. The exchange of pancreas between States and New Zealand in 2004-2006

Source: ANZOD Registry Report 2007

## Bibliography

Australia and New Zealand Organ Donation Registry. ANZOD Registry Report 2008. Adelaide. South Australia. Ed, Herbertt, K. & Russ, G. R.

Australia and New Zealand Organ Donation Registry. ANZOD Registry Report 2007. Appendix I. Adelaide. South Australia. Ed, Herbertt, K. & Russ, G. R. (<http://www.anzdata.org.au/anzod/ANZODReport/2006/2006App1.pdf>)

Australia and New Zealand Organ Donation Registry. Summaries of organ donations and transplants in Australia and New Zealand in 2008. <http://www.organ.redcross.org.au>

Australia and New Zealand Organ Donation Registry. Summaries of organ donations and transplants in Australia and New Zealand in 2007. <http://www.organ.redcross.org.au>

International Pancreas Transplant Registry. Newsletter. Vol. 16(1). August 2004.